10

HANDSMAN & HADDAD PERIODONTICS, P.C.

Practice Limited to Perindonties & Implantology

David R. Handsman, DMD MDS
Todd M. Jenny, DOS, MS
Diplomates, American Board of Periodontology

PRE-OPERATIVE INSTRUCTIONS

- No aspirin 7-10 days prior to surgery. Aspirin and some herbal supplements may cause bleeding. Please check with your physician before stopping aspirin therapy.
- 2. No alcohol for two days prior to surgery.
- One day prior to surgery, begin taking 11buprofen and 1 Extra Strength Tylenol every six hours, and antibiotics as directed.
- Make sure to eat prior to surgery. There is no need to fast. Do not have caffeinated beverages prior to surgery.

POST- OPERATIVE INSTRUCTIONS

- Continue taking medications as directed. Take all antibiotics until finished.
 Continue 1 Ibuprofen and 1 Extra Strength Tylenol every 6 hours for 3-7 days.
- Apply ice pack to surgical site for 15 minutes on and off for the first 24 hours. Use ice day of surgery only.
- Nothing hot to eat or drink the first day. Good nutrition is important during healing. Keep to a softer diet for the week.
- If bleeding occurs apply a moist black tea bag with pressure to the area. It is normal to find blood in the saliva for the first 24 - 48 hours.
- Avoid strenuous physical activity or exertion for 1-2 weeks after surgery.
- 7. Use two pillows when you lie down. Get plenty of rest.
- Do not brush the area of surgery. Use Chlorhexidine rinse 2 times a
 day beginning the day after surgery. Try to keep your mouth as clean as possible
 to promote healing. Chlorhexidine can cause temporary staining with prolonged use.
- If you have any questions or problems (vomiting, rash, hives, itching, or loss of dressing) do not hesitate to call.
- 10. No rinsing, spitting or using a straw for 24 hours.
- It is helpful to take 2 tablespoons of Stonyfield organic yogurt ½ hour before Medications or use probiotics to offset potential stomach problems due to medications.